

## DOCK8 RABBIT MAB

**Cat.#:** N262146

**Product Name:** Anti-DOCK8 Rabbit Monoclonal Antibody

**Synonyms:** MRD2; ZIR8; HEL-205

**UNIPROT ID:** Q8NF50

**Background:** Guanine nucleotide exchange factor (GEF) which specifically activates small GTPase CDC42 by exchanging bound GDP for free GTP (PubMed:28028151, PubMed:22461490). During immune responses, required for interstitial dendritic cell (DC) migration by locally activating CDC42 at the leading edge membrane of DC. Required for CD4+ T-cell migration in response to chemokine stimulation by promoting CDC42 activation at T cell leading edge membrane (PubMed:28028151). Is involved in NK cell cytotoxicity by controlling polarization of microtubule-organizing center (MTOC), and possibly regulating CCDC88B-mediated lytic granule transport to MTOC during cell killing (PubMed:25762780).

**Immunogen:** Recombinant protein of human DOCK8

**Applications:** WB, ICC/IF

**Recommended Dilutions:** WB: 1/500-1/1000 IF: 1/50-1/200

**Host Species:** Rabbit

**Clonality:** Rabbit Monoclonal

**Clone ID:** R05-7D3

**MW:** Calculated MW: 239 kDa; Observed MW: 239 kDa

**Isotype:** IgG

**Purification:** Affinity Purified

**Species Reactivity:** Human


**Conjugation:** Unconjugated

**Modification:** Unmodified

**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

**Research Areas:** Cardiovascular

**Storage & Shipping:** Store at -20°C. Avoid repeated freezing and thawing



Western blot analysis of DOCK8 in Ramos lysates using DOCK8 antibody.