

PI3 KINASE P110 DELTA RABBIT PAB

Cat.#: N225115

Product Name: Anti-PI3 Kinase p110 delta Rabbit pAb

Synonyms: PIK3CD; Phosphatidylinositol 4; 5-bisphosphate 3-kinase catalytic subunit delta isoform; PI3-kinase subunit delta; PI3K-delta; PI3Kdelta; PtdIns-3-kinase subunit delta; Phosphatidylinositol 4; 5-bisphosphate 3-kinase 110 kDa catalytic subunit delta; PtdIns-3-kinase subunit p110-delta; p110delta

UNIPROT ID: O00329

Background: Plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Mediates immune responses. Plays a role in B-cell development, proliferation, migration, and function. Required for B-cell receptor (BCR) signaling.

Immunogen: The antiserum was produced against synthesized peptide derived from the N-terminal region of human PIK3CD. AA range:41-90

Applications: WB,IHC-P,ELISA

Recommended Dilutions: WB: 1/500-1/1000 IHC: 1/50-1/100 ELISA: 1/10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Clone ID: -

MW: Calculated MW: 119 kDa; Observed MW: 119 kDa

Isotype: IgG

Purification: Affinity Purified

Species Reactivity: Human,Mouse


Conjugation: Unconjugated

Modification: Unmodified

Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

Research Areas: Signal Transduction

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin-embedded mouse brain using PI3 Kinase p110 delta antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of PI3 Kinase p110 delta in HepG2 lysates using PI3 Kinase p110 delta antibody.