

PPAR GAMMA RABBIT PAB

Cat.#: N225414

Product Name: Anti-PPAR gamma Rabbit pAb

Synonyms: CIMTI; GLMI; NR1C3; Nuclear receptor subfamily 1 group C member 3; Peroxisome proliferator activated nuclear receptor gamma variant 1; Peroxisome proliferator activated receptor gamma 1; Peroxisome Proliferator Activated Receptor gamma; Peroxisome proliferator-activated receptor gamma; PPAR gamma; PPAR-gamma; PPARG; PPARG_HUMAN; PPARG1; PPARG2; PPARgamma.

UNIPROT ID: P37231

Background: Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the receptor binds to a promoter element in the gene for acyl-CoA oxidase and activates its transcription. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Key regulator of adipocyte differentiation and glucose homeostasis.

Immunogen: Recombinant Protein

Applications: WB,IHC-P

Recommended Dilutions: WB: 1/500-1/1000 IHC: 1/50-1/100

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Clone ID: -

MW: Calculated MW: 58 kDa; Observed MW: 53,57 kDa

Isotype: IgG

Purification: Affinity Purified

Species Reactivity: Human,Mouse,Rat


Conjugation: Unconjugated

Modification: Unmodified


Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

Research Areas: Epigenetics and Nuclear Signaling

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing



Western blot analysis of PPAR gamma in Hela, 3T3, PC-12 lysates using PPAR gamma antibody



Immunohistochemistry analysis of paraffin-embedded mouse Heart Tissue using PPAR gamma antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.