

## RPA32 RABBIT MAB

**Cat.#:** N262844

**Product Name:** Anti-RPA32 Rabbit Monoclonal Antibody

**Synonyms:** 60S acidic ribosomal protein P1; AA409079; AI325195; AU020965; HSSB; ik:tdsubc\_2g1; M(2)21C; MGC137236; OTTHUMP00000004008; p32; p34; RCJMB04\_6d17 replication protein A2; 32kDa; REPA 2; REPA1; REPA2; Replication factor A protein 2; Replication protein A 32 kDa subunit; Replication protein A 32kDa subunit; Replication protein A 34 kDa subunit; Replication protein A; replication protein A1 (70kD); Replication Protein A2 (32kDa); Replication protein A2 32kD; Replication protein A2 32kDa; Replication protein A2; Replication protein A2; 32kDa; RF A; RF-A protein 2; Rf-A2; RFA; RFA2\_HUMAN; RP A; RP-A p32; RP-A p34; RP21C; RPA 2; RPA 32; RPA; RPA2; RPA32; RPA34; RPA70; RplP1; RpP2; xx:tdsubc\_2g1; zgc:109822.

**UNIPROT ID:** P15927

**Background:** As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage. In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation.

**Immunogen:** A synthetic peptide of human RPA32/RPA2

**Applications:** WB,IHC-P,IP

**Recommended Dilutions:** WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20

**Host Species:** Rabbit

**Clonality:** Rabbit Monoclonal

**Clone ID:** R07-4D1

**MW:** Calculated MW: 29 kDa; Observed MW: 29 kDa

**Isotype:** IgG

**Purification:** Affinity Purified

**Species Reactivity:** Human,Mouse,Rat,Hamster


**Conjugation:** Unconjugated

**Modification:** Unmodified


**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

**Research Areas:** Epigenetics and Nuclear Signaling


**Storage & Shipping:** Store at  $-20^{\circ}\text{C}$ . Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin-embedded Human tonsil using RPA32/RPA2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of RPA32/RPA2 in 3T3 lysates using RPA32 antibody.



Western blot analysis of RPA32/RPA2 in Hela, CHO-K1, C6, Jurkat lysates using RPA32/RPA2 antibody.