

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

ZAP70 RABBIT PAB

Cat.#: S219146

Product Name: Anti-ZAP70 Rabbit Polyclonal Antibody **Synonyms:** SRK; STD; TZK; STCD; IMD48; ADMIO2; ZAP-70 UNIPROT ID: P43403 (Gene Accession - BC039039)

Background: This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it

plays a role in T-cell development and lymphocyte activation. This enzyme, which is

phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been

found for this gene.

Immunogen: Fusion protein of human ZAP70

Applications: ELISA, WB, IHC

Recommended Dilutions: IHC: 50-300;WB: 500-2000;ELISA: 5000-10000

Host Species: Rabbit

Clonality: Rabbit Polyclonal

Isotype: Immunogen-specific rabbit IgG **Purification:** Antigen affinity purification Species Reactivity: Human, Mouse

Constituents: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40%

glycerol


Research Areas: Signal Transduction, Immunology

Storage & Shipping: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 219146(ZAP70 Antibody) at a dilution of 1/55(Cytoplasm).



In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the fusion protein and then with 219146 (Anti-ZAP70 Antibody) at dilution 1/55.



Gel: 8%SDS-PAGE, Lysate: 40 µg;

Lane 1-2: Mouse liver tissue and Jurkat cell lysates;

Primary antibody: 219146(ZAP70 Antibody) at

dilution 1/800;

Secondary antibody: HRP-conjugated Goat

anti rabbit IgG at 1/5000 dilution;

Exposure time: 3 minutes