

ATM PAB

ATM Polyclonal Antibody

Cat. #: 21276

Gene Symbol: ATM

Description: Anti-ATM Rabbit Polyclonal Antibody

Background: Ataxia telangiectasia mutated (ATM) is a serine/threonine protein kinase that is recruited and activated by DNA double-strand breaks. It phosphorylates several key proteins that initiate activation of the DNA damage checkpoint, leading to cell cycle arrest, DNA repair or apoptosis. Activity of ATM protein is under tight control, and mutation of ATM can cause disease such as Ataxia telangiectasia (AT) and cancers.

Immunogen: Purified ATM protein, human origin.

Applications: ELISA, WB, IHC

Recommended Dilutions:

ELISA: 1:1000-1:5000

WB: 1:100-1:1000

IHC: 1:50-1:200

Concentration: 0.2 mg/ml

Host Species: Rabbit

Format: Liquid

Clonality: Polyclonal

Isotype:

Purity: Purified from serum

Preservative: No


Constituents: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 50% glycerol

Species Reactivity: Recognizes ATM of vertebrates.

Storage Conditions: Store at -20°C. Avoid repeated freezing and thawing

Western blot:


ATM protein



WB: Anti-ATMpAb

Western blot analysis of recombinant ATM protein. Purified His-tagged ATM protein was blotted with Anti-ATM rabbit polyclonal antibody (Cat. #21276).

Immunohistochemistry:



Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue with Anti-ATM polyclonal antibody (Cat. #21276). Tissue samples were fixed with formaldehyde and blocked with 1% serum for 15 min at 37 °C. Antigen retrieval was by heat mediation in citrate buffer (pH6). Samples were then incubated with primary antibody (1:50 dilution) overnight at 4°C. A HRP-conjugated Goat anti-mouse IgG (1:50 dilution) was used as secondary antibody.