

CCR1 (DMC465) IGG1 CHIMERIC MAB

目录: 28241

产品名称: CCR1(DMC465) IgG1 Chimeric Monoclonal Antibody

基因符号: CD191; CKR-1; CKR1; CMKBR1; HMI45; MIP1aR; SCYAR1

描述: CCR1 antibody(DMC465) IgG1 Chimeric Monoclonal Antibody

背景: This gene encodes a member of the beta chemokine receptor family; which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. The ligands of this receptor include macrophage inflammatory protein 1 alpha (MIP-1 alpha); regulated on activation normal T expressed and secreted protein (RANTES); monocyte chemoattractant protein 3 (MCP-3); and myeloid progenitor inhibitory factor-1 (MPIF-1). Chemokines and their receptors mediated signal transduction are critical for the recruitment of effector immune cells to the site of inflammation. Knockout studies of the mouse homolog suggested the roles of this gene in host protection from inflammatory response; and susceptibility to virus and parasite. This gene and other chemokine receptor genes; including CCR2; CCRL2; CCR3; CCR5 and CXCR1; are found to form a gene cluster on chromosome 3p. [provided by RefSeq; Jul 2008]

经过测试的应用: Flow Cyt

推荐稀释度: Flow Cyt 1:100

种属反应性: Rabbit

亚型: Rabbit:Human Fc chimeric IgG1

纯化: Purified from cell culture supernatant by affinity chromatography

种属反应性: Human CCR1

成分: Lyophilized from sterile PBS, pH 7.4. 5 % – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).




Figure 1. CCR1 protein is highly expressed on the surface of Expi293 cell membrane. Flow cytometry analysis with CCR1 (DMC465) on Expi293 cells transfected with human CCR1 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram), and Isotype antibody on Expi293 transfected with irrelevant protein (Orange histogram).