

CD22 (DM13) RABBIT MAB

目录: 28301

产品名称: CD22(DM13) Rabbit Monoclonal Antibody

基因符号: SIGLEC-2; SIGLEC2

描述: CD22 antibody(DM13) Rabbit Monoclonal Antibody

背景: CD22 (CD22 Molecule) is a Protein Coding gene. Diseases associated with CD22 include Refractory Hematologic Cancer and Hairy Cell Leukemia. Among its related pathways are Downstream signaling events of B Cell Receptor (BCR) and Hematopoietic cell lineage. Gene Ontology (GO) annotations related to this gene include carbohydrate binding. An important paralog of this gene is SIGLEC2.

经过测试的应用: ELISA; Flow Cyt

推荐稀释度: Flow Cyt 1:100

种属反应性: Rabbit

亚型: Rabbit IgG

纯化: Purified from cell culture supernatant by affinity chromatography

种属反应性: Human CD22

成分: Lyophilized from sterile PBS, pH 7.4. 5 % - 8 % trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).




Figure 1. Expi 293 cell line transfected with irrelevant protein (left) and human CD22 (right) were surface stained with Rabbit CD22 monoclonal antibody 1µg/ml (clone: DM13) followed by Alexa 488-conjugated rabbit IgG secondary antibody.




Figure 2. Flow cytometry data of serially titrated Rabbit CD22 monoclonal antibody (clone: DM13) on Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.




Figure 3. Affinity ranking of different Rabbit CD22 mAb clones by titration of different concentration onto Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.




Figure 4. Phylogenetic analysis of amino acid sequence of different Rabbit CD22 mAb clones. A) Heavy chain and B) Light chain.