

**HUMAN AGR2 PROTEIN, HFC TAG**

目录: 11571

产品名称: Human AGR2 Protein

规格: 10 µg, 50 µg and 100 µg

基因符号: AG-2;hAG-2;HPC8

**Target:** AGR2

**UNIPROT ID:** O95994

**描述:** Recombinant human AGR2 protein with C-terminal human Fc tag

**背景:** This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, a catalytically active thioredoxin domain, and a C-terminal ER-retention sequence. This protein plays a role in cell migration, cellular transformation and metastasis and is as a p53 inhibitor. As an ER-localized molecular chaperone, it plays a role in the folding, trafficking, and assembly of cysteine-rich transmembrane receptors and the cysteine-rich intestinal glycoprotein mucin. This gene has been implicated in inflammatory bowel disease and cancer progression. [provided by RefSeq, Mar 2017]

**Species/Host:** HEK293

**Molecular Weight:** The protein has a predicted molecular mass of 44.0 kDa after removal of the signal peptide. The apparent molecular mass of AGR2-hFc is approximately 35-55 kDa due to glycosylation.

**Molecular Characterization:** AGR2(Arg21-Leu175) hFc(Glu99-Ala330)

**纯化::** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

**Formulation & Reconstitution:** Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

**储存和运输:** Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human AGR2 Protein, hFc Tag on SDS-PAGE under reducing condition.