

HUMAN BCMA PROTEIN, HIS TAG

目录: 11314

产品名称: Human BCMA Protein

规格: 10 µg, 50 µg and 100 µg

基因符号: TNFRSF17;CD269;BCM;BCMA

Target: BCMA

UNIPROT ID: Q02223

描述: Recombinant Human BCMA Protein with C-terminal 6xHis tag

背景: B-cell maturation protein (BCMA or BCM), is also known as Tumor necrosis factor receptor superfamily member 17 (TNFRSF17), which is encoded by the TNFRSF17 gene. TNFRSF17 is a cell surface receptor of the TNF receptor superfamily which recognizes B-cell activating factor (BAFF). This receptor is expressed in immune organs and mature B cell lines. BCMA promotes B-cell survival and plays a role in the regulation of humoral immunity. BCMA can activate NF-kappa-B and JNK.

Species/Host: HEK293

Molecular Weight: The protein has a predicted molecular mass of 6.7kDa after removal of the signal peptide.

Molecular Characterization: BCMA(Met1-Ala54) 6×His tag

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human BCMA Protein, His Tag on SDS-PAGE under reducing condition.

Human BCMA, His Tagged protein ELISA

0.2 µg of BCMA, His Tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human BAFF, hFc tagged protein 11169 can bind Human BCMA, His tagged protein (11314) in a linear range of 1.28-160 ng/ml.

