

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **HUMAN CD112 PROTEIN, HFC TAG**

目录: 11353

产品名称: Human CD112 Protein 规格: 10 µg, 50 µg and 100 µg

基因符号: NECTIN2; HVEB; PRR2; PVRL2; PVRR2

Target: CD112

**UNIPROT ID:** Q92692

描述: Recombinant Human CD112 Protein with C-terminal human Fc

tag

背景: This gene encodes a single-pass type I membrane glycoprotein with two Ig-like C2-type domains and an Ig-like V-type domain. This protein is one of the plasma membrane components of adherens junctions. It also serves as an entry for certain mutant strains of herpes simplex virus and pseudorabies virus, and it is involved in cell to cell spreading of these viruses. Variations in this gene have been associated with differences in the severity of multiple sclerosis. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

Species/Host: HEK293

**Molecular Weight:** The protein has a predicted molecular mass of 61.4 kDa after removal of the signal peptide. The apparent molecular mass of CD112-hFc is approximately 70 kDa due to glycosylation.

Molecular Characterization: CD112(Gln32-Gly360) hFc(Glu99-Ala330)

纯化:: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

**Formulation & Reconstitution:** Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010




Figure 1. Human CD112 Protein, hFc Tag on SDS-PAGE under reducing condition.




Figure 2. Flow cytometry analysis with 15µg/ml Human CD112 Protein, hFc tag (11353) on Expi293 cells transfected with human PVRIG (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).