

IDH2(R172K)**IDH2(R172K)**

货号: 26163

基因符号: IDH2; D2HGA2; ICD-M; IDH; IDHM; IDP; IDPM; mNADP-IDH

描述: IDH2(R172K) 小鼠单克隆抗体

背景: Isocitrate dehydrogenase (IDH) catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. The isocitrate and isopropylmalate dehydrogenases family has three members, IDH1, IDH2 and IDH3. IDH2 plays a role in intermediary metabolism and energy production. Defects in IDH2 are the cause of D-2-hydroxyglutaric aciduria type 2 (D2HGA2). Somatic mosaic mutations of this protein have also been found associated to Ollier disease and Maffucci syndrome, and R172K IDH2 mutations do exist in diffusely infiltrative gliomas.

免疫原: A synthetic peptide from the internal region of IDH2 which includes the mutation of R172K, human origin.

经过测试的应用: ELISA, WB, IHC

推荐稀释度:

ELISA: 1:1000-1:5000

WB: 1:100-1:1000

IHC: 1:50-1:100

IHC: 1:10-1:100

浓度: 1 mg/ml

种属反应性: Mouse

形式: Liquid

克隆性: Monoclonal

亚型:

纯化: Purified from ascites

Preservative: No

成分: PBS (without Mg^{2+} and Ca^{2+}), pH 7.4, 150 mM NaCl, 50% glycerol


种属反应性: recognizes R172K mutant, but not wild type IDH2 of vertebrates.

储存条件: Store at $-20^{\circ}C$. Avoid repeated freezing and thawing

Western blot:

IDH2 protein

WT R172K




1 2


WB: anti-IDH2(R172K) mAb

Western blot analysis of recombinant IDH2(R172K) and wild type proteins.

Purified His-tagged IDH2(R172K) (lane 2) and wild type protein (lane 1) were blotted with IDH2(R172K) monoclonal antibody (货号26163).

Immunofluorescence:





Immunofluorescence of cells expressing IDH2 proteins with IDH2(R172K) antibody.

HEK293T cells were transfected with pCDNA3-GFP- IDH2(R172K) plasmid , pCDNA3-GFP- IDH2 (WT) plasmid, pCDNA3-GFP- IDH2 (R172H) plasmid , pCDNA3-GFP- IDH2 (R172W) plasmid, pCDNA3-GFP- IDH2 (R172G) plasmid,pCDNA3-GFP- IDH2 (R172M) plasmid or pCDNA3-GFP- IDH2 (R172S) plasmid, then fixed and stained with IDH2(R172K) monoclonal antibody (货号26163).