

## **NECTIN4 (DMC438) IGG1 CHIMERIC MAB**

目录: 28226

产品名称: Nectin4(DMC438) IgG1 Chimeric Monoclonal Antibody

基因符号: EDSS1; LNIR; nectin-4; PRR4; PVRL4

描述: Nectin4 antibody(DMC438) IgG1 Chimeric Monoclonal Antibody

背景: This gene encodes a member of the nectin family. The encoded protein contains two immunoglobulin-like (Ig-like) C2-type domains and one Ig-like V-type domain. It is involved in cell adhesion through trans-homophilic and -heterophilic interactions. It is a single-pass type I membrane protein. The soluble form is produced by proteolytic cleavage at the cell surface by the metalloproteinase ADAM17:TACE. The secreted form is found in both breast tumor cell lines and breast tumor patients. Mutations in this gene are the cause of ectodermal dysplasia-syndactyly syndrome type 1; an autosomal recessive disorder. Alternatively spliced transcript variants have been found but the full-length nature of the variant has not been determined.

经过测试的应用: Flow Cyt

推荐稀释度: Flow Cyt 1:100

种属反应性: Rabbit

亚型: Rabbit:Human Fc chimeric IgG1

纯化: Purified from cell culture supernatant by affinity chromatography

种属反应性: Human NECTIN4

成分: Lyophilized from sterile PBS, pH 7.4. 5 % – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form.

After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).




Figure 1. Flow cytometry analysis with Nectin4 (DMC438) on Expi293 cells transfected with human Nectin4 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).