

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

HUMAN TLR2 FULL LENGTH PROTEIN

目录: 11121 产品名称: Human TLR2 Full Length Protein 规格: 10 µg, 50 µg and 100 µg 基因符号: CD282; TIL4

Target: TLR2

UNIPROT ID: 060603

描述: Human TLR2 full length protein-synthetic nanodisc

背景: The protein is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. This protein is a cell-surface protein that can form heterodimers with other TLR family members to recognize conserved molecules derived from microorganisms known as pathogen-associated molecular patterns (PAMPs). Activation of TLRs by PAMPs leads to an up-regulation of signaling pathways to modulate the host's inflammatory response. This gene is also thought to promote apoptosis in response to bacterial lipoproteins. This gene has been implicated in the pathogenesis of several autoimmune diseases. Alternative splicing results in multiple transcript variants.

Species/Host: HEK293


Molecular Weight: The human full length TLR2 protein has a MW of 89.8 kDa **Formulation & Reconstitution:** Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Product Description

Pioneering GTPase and Oncogene Product Development since 2010



Figurel. Elisa plates were pre-coated with Flag Tag TLR2-Nanodisc (0.2 µg/per well). Serial diluted Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for Flag monoclonal antibody binding with TLR2-Nanodisc is 2.685ng/ml.




Figure2. Human TLR2-Nanodisc, Flag Tag on SDS-PAGE