

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

IDH2(R172W)

Catalog Number: 26164

Gene Symbol: IDH2; D2HGA2; ICD-M; IDH; IDHM; IDP; IDPM; mNADP-IDH

Description: Anti-IDH2(R172W) Mouse Monoclonal Antibody

Background: Isocitrate dehydrogenase (IDH) catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. The isocitrate and isopropylmalate dehydrogenases family has three members, IDH1, IDH2 and IDH3. IDH2 plays a role in intermediary metabolism and energy production. Defects in IDH2 are the cause of D-2-hydroxyglutaric aciduria type 2 (D2HGA2). Somatic mosaic mutations of this protein have also been found associated to Ollier disease and Maffucci syndrome, and R172W IDH2 mutations do exist in diffusely infiltrative gliomas.

Immunogen: A synthetic peptide from the internal region of IDH2, which includes the mutation of

R172W, human origin.

Tested Applications: ELISA, WB, IF, IHC

Recommended Dilutions:

ELISA: 1:1000-1:5000 WB: 1:500-1:1000 IF: 1:50-1:100 IHC: 1:50-1:100

Concentration: 1 mg/ml

Host: Mouse

Clonality: Monoclonal

Isotype: IgG

Purity: Purified from ascites

Format: Liquid Preservative: No

Constituents: PBS (without Mg²⁺ and Ca²⁺), pH7.4, 150

mM NaCl, 50% glycerol


Species Reactivity: Recognizes R172W mutant, but not

wild-type IDH2 of vertebrates.

Storage Conditions: Store at -20°C. Avoid repeated

freezing and thawing

Western blot:



WB: Anti-IDH2(R172W) mAb

Western blot analysis of recombinant IDH2 (R172W) and wild type proteins.

Purified His-tagged IDH2 (R172W) (lane 2) and wild type protein (lane 1) were blotted with anti-IDH2 (R172W) monoclonal antibody (Cat. # 26164).

NewEast Biosciences 1150 First Avenue, Suite 501 King of Prussia, PA 19406 Support:

610-945-2007


info@neweastbio.com

Web: www.neweastbio.com



Pioneering GTPase and Oncogene Product Development since 2010

Immunofluorescence:



Immunofluorescence of cells expressing IDH2 proteins with anti-IDH2 (R172W) antibody.

HEK293T cells were transfected with pCDNA3-GFP-IDH2 (WT) plasmid (left column) or pCDNA3-GFP-IDH2 (R172W) plasmid (right column), then fixed and stained with anti-IDH2 (R172W) monoclonal antibody (Cat. # 26164).

Web: www.neweastbio.com